

The new intermediate long bursting source XTE J1701-407

M. Falanga,^{1,2} A. Cumming,³ E. Bozzo,^{4,5} J. Chenevez⁶

¹ CEA Saclay, DSM/IRFU/Service d'Astrophysique (CNRS FRE 2591), 91191, Gif sur Yvette, France e-mail: mfalanga@cea.fr

² AIM - Unité Mixte de Recherche CEA - CNRS - Université Paris 7, Paris, France

³ Physics Department, McGill University, 3600 rue University, Montreal QC, H3A 2T8, Canada

⁴ INAF - Osservatorio Astronomico di Roma, Via Frascati 33, 00044 Rome, Italy

⁵ Dipartimento di Fisica - Università di Roma Tor Vergata, via della Ricerca Scientifica 1, 00133 Rome, Italy

⁶ National Space Institute, Technical University of Denmark, Juliane Maries Vej 30, 2100 Copenhagen, Denmark

Preprint online version: January 22, 2009

ABSTRACT

Aims. XTE J1701-407 is a newly discovered X-ray transient source. In this work we investigate its flux variability and study the intermediate long and short bursts discovered by *Swift* on July 17, and 27, 2008, respectively.

Methods. So far, only one intermediate long burst, with a duration of ≈ 18 minutes and ten days later a short burst, have been recorded from XTE J1701-407. We analyzed the public available data from *Swift* and *RXTE*, and compared the observed properties of the intermediate long burst with theoretical ignition condition and light curves to investigate the possible nuclear burning processes.

Results. The intermediate long burst may have exhibited a photospheric radius expansion, allowing us to derive the source distance at 6.2 kpc assuming the empirically derived Eddington luminosity for pure helium. The intermediate long burst decay was best fit by using two exponential functions with e-folding times of $\tau_1 = 40 \pm 3$ s and $\tau_2 = 221 \pm 9$ s. The bursts occurred at a persistent luminosity of $L_{\text{per}} = 8.3 \times 10^{36}$ erg s⁻¹ ($\approx 2.2\%$ of the Eddington luminosity). For the intermediate long burst the mass accretion rate per unit area onto the NS was $\dot{m} \approx 4 \times 10^3$ g cm⁻² s⁻¹, and the total energy released was $E_{\text{burst}} \approx 3.5 \times 10^{40}$ erg. This corresponds to an ignition column depth of $y_{\text{ign}} \approx 1.8 \times 10^9$ g cm⁻², for a pure helium burning. We find that the energetics of this burst can be modeled in different ways, as (i) pure helium ignition, as the result of either pure helium accretion or depletion of hydrogen by steady burning during accumulation, or (ii) as ignition of a thick layer of hydrogen-rich material in a source with low metallicity. However, comparison of the burst duration with model light curves suggests that hydrogen burning plays a role during the burst, and therefore this source is a low accretion rate burster with a low metallicity in the accreted material.

Key words. binaries: close – stars: individual: XTE J1701-407 – stars: neutron – X-rays: bursts

1. Introduction

Type I X-ray bursts are among the most evident signatures that testify the presence of a neutron star (NS) in low mass X-ray binaries. These bursts are thermonuclear explosions that occur on the surface of accreting NSs and are triggered by unstable hydrogen and/or helium burning (see, e.g., Lewin, van Paradijs & Taam, 1993; Strohmayer & Bildsten, 2006, for reviews). Type I X-ray bursts were predicted theoretically by Hansen & Van Horn (1975), and several thousand bursts have been observed to date (see, e.g., Cornelisse et al., 2003; Galloway et al., 2006a; Chelovekov, Grebenev & Sunyaev, 2006). From the duration of the bursts measured by their decay parameter τ (see e.g., Galloway et al., 2006a) three main branches are distinguished: normal bursts, intermediate long bursts, and superbursts (see Fig. 7 in Falanga et al., 2008, and references therein). These bursts can be described by different fuel types, accretion rates, and they, therefore, also displays different recurrence times (e.g., Strohmayer & Bildsten, 2006; Cumming & Macbeth, 2004; Galloway et al., 2006a). Thanks to the long Galactic plane scan carried out with *BeppoSAX*, *INTEGRAL*, and *Swift* a number of rare inter-

mediate long bursts and superbursts have been observed. In most cases the rise of the burst is of ≈ 1 s, whereas the decay is approximately exponential, with a duration of a few seconds for normal bursts, tens of minutes for intermediate long bursts, and up to several hours for superbursts (e.g., Kuulkers, 2004; in 't Zand et al., 2004; Molokov et al., 2005; in 't Zand et al., 2005; Chenevez et al., 2006, 2007; Falanga et al., 2008). The recurrence time of type I X-ray bursts ranges from few hours to years, depending on the nuclear reactions involved (see, e.g., Lewin, van Paradijs & Taam, 1993; Strohmayer & Bildsten, 2006, for reviews).

In this paper, we report on an intermediate long burst from the X-ray transient XTE J1701-407. This source was discovered by the *Rossi X-ray Timing Explorer (RXTE)* during a routine Galactic bulge scan on June 8, 2008 (Markwardt et al., 2008a). At the time of the discovery, the source spectrum was best fit with an absorbed power-law model with photon index ≈ 2.2 and a neutral absorption column of $\approx 3.4 \times 10^{22}$ cm⁻², the measured flux was $\approx 1.4 \times 10^{-10}$ erg cm⁻² s⁻¹ in the 2-10 keV band (Markwardt et al., 2008a). *Swift*/XTE follow-up observations on June 11, 2008 found both spectral and source flux measurements consistent with the *RXTE*/PCA re-




Fig. 1. Flux history of the X-ray transient XTE J1701-407. Open squares represent fluxes derived from *RXTE* bulge observations in the period from March 31, to September 2, 2008. Fluxes derived from *RXTE*/PCA and *Swift*/XRT target of opportunity and follow-up observations are marked with filled squares and stars, respectively. The arrow indicates the time of the type I X-ray burst.

sults (Degenaar & Wijnands, 2008). On July 17, 2008, the *Swift*/BAT camera detected a short flare consistent with the position of XTE J1701-407, and ~ 97 s later *Swift*/XRT measured a decaying X-ray flux (Barthelmy et al., 2008). Based on this BAT data the X-ray spectrum was consistent with those observed for thermonuclear X-ray bursts, therefore, this flare was associated to a most likely type I burst (Markwardt et al., 2008b). Moreover, based on the *Swift*/XRT time resolved spectrum during the flux decay, Linares et al. (2008a) measured the cooling tail and confirmed the flare to be an intermediate long type I X-ray burst. This led to the classification of XTE J1701-407 as a NS low-mass X-ray binary (Linares et al., 2008a,b). Ten days later, i.e., on July 27, BAT observed in the 15–150 keV band a short bursts lasting ≈ 10 s (Sakamoto et al., 2008). Kilohertz quasi-periodic oscillations were observed at ≈ 1150 Hz (see Strohmayer, Markwardt & Swank, 2008, for details). The most accurate position of the source was provided by *Swift* at $\alpha_{J2000}=17^{\text{h}}01^{\text{m}}44^{\text{s}}.3$ and $\delta_{J2000}=-40^{\circ}51'29''.9$ with an estimated accuracy of $1''.7$ (Starling & Evans, 2008). No infrared counterpart candidate was found at this position (Kaplan & Chakrabarty, 2008).

2. Data Analysis and results

For the present study we used public available data from *Swift* and *RXTE* observatories. The dataset include *RXTE* target of opportunity observations, performed after the discovery of the source (observation ID. 93444), as well as *Swift* follow-up observations (ID. 0031221001, 0031221002, and 00317205001). The *Swift* data set includes also the intermediate long burst discovered on July 17, (ID. 00317205000) and the short burst from July 27, 2008 (ID. 00318166000). The total effective exposure time is 75.4 ks (30 pointings) and 1.1 ks for *RXTE*/PCA and *Swift* respectively. In Table 1 we report the detailed observations log. Data reduction of the *RXTE* Proportional Counter Array (PCA; 2–60 keV, Jahoda et al., 1996) and the High Energy X-ray Timing Experiment (HEXTE; 15–250 keV, Rothschild et al., 1998) was performed according to the

default selection criteria for background subtraction, light curve, and spectrum extraction. We used only PCU2 data. We used *Swift*/XRT data in both Windowed Timing (WT) and Photon Counting (PC) mode. However, in the following we report only on WT data, since we found that PC data were strongly affected by pile-up and quasi-simultaneous observations carried out in WT mode were available. We reduced all the XRT data by using the version of the *xrtpipeline* (version 0.11.6) included in the *Heasoft* package 6.4, and the latest calibration files available. Source events in WT mode were extracted from rectangle regions with widths of 40 pixels and heights of 20 pixels¹. Ancillary response files were generated with *xrtmkarf* and accounted for different extraction regions, vignetting, and point-spread function. For the *Swift*/BAT data analysis we used the *batgrbproduct* tool included in the *Heasoft* package 6.4. Time resolved spectra and lightcurves (see Sec. 2.2) were extracted by using the *batbinevt*, *batupdatephakw*, *batphasyserr*, and *batdrngen* tools.

2.1. Persistent emission

Spectra obtained from *RXTE*/PCA and *Swift*/XRT observations were fit separately by using a photoelectrically-absorbed power-law model. The best fit parameters for the absorption column, N_{H} , are between 1.7 ± 1 and $4.4 \pm 1.2 \times 10^{22} \text{ cm}^{-2}$ and a power-law index $\Gamma = 2.0 \pm 0.1 - 2.7 \pm 0.3$ the corresponding unabsorbed fluxes are between $1.1 \pm 0.07 \times 10^{-10}$ and $8.3 \pm 0.6 \times 10^{-10}$ in the 2–20 keV energy band. In Table 1 we report all the persistent spectral parameters. All uncertainties are at a 90% confidence level. PCA spectra were extracted in the 2–20 keV energy band; the source was only detected at low significance in the HEXTE (15–50 keV band). A fit to the joined broadband PCA/HEXTE (2–50 keV) spectrum did not lead to a significant improvement in the determination of the model parameters; therefore, we report in the following only on the PCA data. *Swift*/XRT spectra were extracted in the 2–10 keV band. All the measured unabsorbed fluxes were extrapolated to the 0.3–100 keV band by generating dummy responses (XSPEC version 11.3.2ag), and are shown in Fig. 1. We included in this figure also fluxes derived from *RXTE* bulge observations² (Swank & Markwardt, 2001). In these cases, the conversion between *RXTE* count rate and flux was obtained using the spectra results and the values reported by Markwardt et al. (2008a).

The light curve and spectral analysis shows that XTE J1701-407 underwent flaring activity lasting a few days, during which the X-ray flux has varied by over a factor of 2–3. This variability can be ascribed to changes in the mass accretion rate or in the position of the source along its colour-colour diagram (CCD, see e.g., Hasinger & van der Klis, 1989). The latter possibility is investigated in Fig. 2. The CCD was realized by using background subtracted *RXTE*/PCA light curves with a 516 s time resolution. The soft colour is defined as the logarithm of the ratio between count rates in the energy bands 2.1–3.7 keV and 3.7–5.7 keV, whereas for the hard colour the energy bands 5.7–9.8 keV and 9.8–18.9 keV are used. No

¹ See also the XRT analysis manual at http://swift.gsfc.nasa.gov/docs/swift/analysis/xrt_swguide_v1.2.pdf

² <http://lheawww.gsfc.nasa.gov/users/craigm/galscan/main.html>

obvious transition between a hard to a soft state is observed. In order to reduce the errors on the colours, we generated a hard intensity diagram (HID) based on the net count rates in the 2.1–5.7 keV and 5.7–18.9 keV with a 516 s time bin. However, the source behaviour is not much better traced in the HID, because the statistical uncertainties are not significantly reduced along the two axes. In Fig. 2 we show the HID. Future observations are thus required in order to investigate further the origin of flux variations in XTE J1701-407.




Fig. 2. HID of XTE J1701-407. The hardness is the ratio of the count rates in the 5.7–18.9 keV to 2.1–5.7 keV and the intensity is in the 2.1–18.9 keV count rate. Each point corresponds to 516 s of integration time.

2.2. Intermediate long burst light curves and spectra

In Fig. 3, we show the *Swift*/BAT 15–150 keV (upper panel) and XRT 0.3–10 keV (lower panel) light curve of the intermediate long burst. The burst start time, is the time at which the BAT X-ray intensity of the source rose to 10% of the peak above the persistent intensity level. The XRT light curve starts with a delay of 133 s after the beginning of the burst as seen by BAT. The BAT light curve shows a slow rise time³ of ≈ 45 s. The XRT decay time from the burst is best fit by using two exponential functions, with e-folding times of $\tau_1 = 44 \pm 8$ s and $\tau_2 = 271 \pm 21$ s, respectively (see also Linares et al., 2008a). The total duration of the burst, i.e. the time spent to go from and back to the persistent state, was of ≈ 86 s and ≈ 18 minutes in the BAT (15–150 keV) and the XRT (0.3–10 keV) light curves, respectively.

Type I X-ray bursts are produced by unstable burning of accreted matter on the surface of the NS. The emission can be well described by a black-body radiation with temperatures, kT_{bb} , in the energy range of a few keV. The energy dependent decay time of these bursts is attributed to the cooling of the NS photosphere resulting in a gradual softening of the burst spectrum. For a review, see, e.g. Lewin, van Paradijs & Taam (1993); Strohmayer & Bildsten (2006). The time-resolved spectral




Fig. 3. The intermediate long type I X-ray bursts detected from XTE J1701-407 on July 17, 2008. The time T_0 expressed in UTC corresponds to $17^{\text{h}}31^{\text{m}}55^{\text{s}}$. The *Swift*/BAT (15–150 keV) and XRT (0.3–10 keV) light curve are shown with a time bin of 5 s. The data gap of ≈ 133 s in the XRT light curve is due to the time elapsed between the BAT trigger and the follow-up observation. The dashed line indicates the background level measured ≈ 3000 s after the burst end.

analysis of the burst was carried out by using BAT and XRT data in the 15–35 keV and 0.3–10 keV bands, respectively. Burst spectra were well fit by a simple photoelectrically-absorbed black-body, BB, model. For all these spectral fit the N_{H} value was frozen at $3.1 \times 10^{22} \text{ cm}^{-2}$ derived from the pre-burst persistent emission (see Table 1). The inferred BB temperature, kT_{bb} , apparent BB radius at 6.2 kpc (see Sec. 3.1), R_{bb} , and bolometric luminosity are shown in Fig. 5 and in Table 3. The burst fluence is calculated from bolometric fluxes, F_{bol} ; these correspond to the observed 2–10 keV *Swift*/XTR fluxes extrapolated to the 0.1–100 keV energy range. The peak flux, F_{peak} , is derived from the BAT 15–35 keV light curve spectra and extrapolated to the 0.1–100 keV energy range. All the measured unabsorbed fluxes were extrapolated to the 0.1–100 keV band by generating dummy responses (XSPEC version 11.3.2ag). This is well justified for the XRT data since the black-body temperature is well inside the spectra bandpass. However, during the main peak a black-body temperature reaches a maximum between 2–3 KeV (see e.g., Falanga et al., 2008, and references therein), and therefore extrapolating the black-body spectra outside the BAT bandpass. For the BAT (15–35 keV) burst peak black-body spectral best fit we found $kT_{\text{bb}} = 2.68 \pm 0.21$ ($\chi^2 = 0.74$, 7 d.o.f.). We fixed an upper and lower boundary black-body temperature of 2 and 3 keV, respectively, leaving the normalization as a free parameters. Note the lower boundary is also consistent to be higher as the first XRT black-body temperature of $kT_{\text{bb}} \approx 1.8$ (see Table 3). We found an unacceptable fit by using a 2 keV black-body burst peak temperature ($\chi^2_{\text{red}} = 5.1$, 8 d.o.f.), whereas a slightly better fit is obtained by using a temperature of 3 keV ($\chi^2_{\text{red}} = 1.22$, 8 d.o.f.). In Fig. 4 we show the BAT data with the different black-body models from

³ The rise time is defined as the time spent between the start of the burst and the point at which the 90% of the peak burst intensity is reached.

Table 1. Log of *RXTE* and *Swift* observations and best fit spectral parameters of the persistent emission. An absorbed power-law model is used to fit these spectra.

Instrument	Date MJD	Exp. ks	N_H 10^{22} cm^{-2}	Γ	Flux $_{2-10\text{keV}}^a$ $\text{erg cm}^{-2} \text{ s}^{-1}$	Flux $_{2-20\text{keV}}^a$ $\text{erg cm}^{-2} \text{ s}^{-1}$	Flux $_{0.3-100\text{keV}}^a$ $\text{erg cm}^{-2} \text{ s}^{-1}$	χ_{red}^2
<i>Swift</i> ^b	54628.66385	0.4	$4.0^{+0.7}_{-0.6}$	$2.2^{+0.3}_{-0.3}$	$2.8 \pm 0.3\text{e-10}$	$3.7 \pm 0.5\text{e-10}$	$10.2 \pm 2\text{e-10}$	0.8
<i>RXTE</i>	54633.24637	2.0	$3.4^{+1.2}_{-2}$	$2.4^{+0.1}_{-0.2}$	$4.5 \pm 0.4\text{e-10}$	$5.6 \pm 0.4\text{e-10}$	$17.5 \pm 7\text{e-10}$	1.2
<i>RXTE</i>	54634.42452	2.5	$3.4^{+1.2}_{-1.2}$	$2.4^{+0.1}_{-0.2}$	$3.4 \pm 0.2\text{e-10}$	$4.3 \pm 0.2\text{e-10}$	$12.8 \pm 2\text{e-10}$	0.7
<i>RXTE</i>	54635.27545	2.0	$3.3^{+2.1}_{-2.2}$	$2.2^{+0.1}_{-0.2}$	$2.7 \pm 0.2\text{e-10}$	$3.6 \pm 0.2\text{e-10}$	$9.4 \pm 1.6\text{e-10}$	0.8
<i>RXTE</i>	54638.80878	3.5	$3.4^{+0.8}_{-0.9}$	$2.2^{+0.1}_{-0.1}$	$1.9 \pm 0.1\text{e-10}$	$2.6 \pm 0.1\text{e-10}$	$6.8 \pm 1.1\text{e-10}$	0.9
<i>RXTE</i>	54639.85563	3.0	$3.4^{+0.7}_{-0.7}$	$2.3^{+0.1}_{-0.1}$	$2.1 \pm 0.1\text{e-10}$	$2.8 \pm 0.1\text{e-10}$	$7.8 \pm 1.4\text{e-10}$	1.2
<i>RXTE</i>	54640.90230	3.5	$3.5^{+0.7}_{-0.7}$	$2.2^{+0.1}_{-0.1}$	$1.9 \pm 0.1\text{e-10}$	$2.6 \pm 0.1\text{e-10}$	$7.1 \pm 1.1\text{e-10}$	1.2
<i>Swift</i> ^{b,c}	54664.62684	0.6	$3.1^{+0.4}_{-0.4}$	$1.9^{+0.2}_{-0.2}$	$4.9 \pm 0.3\text{e-10}$	$7.2 \pm 0.6\text{e-10}$	$18.3 \pm 3\text{e-10}$	1.1
<i>RXTE</i>	54665.10546	3.0	$3.7^{+1.2}_{-2}$	$2.6^{+0.2}_{-0.1}$	$4.9 \pm 0.6\text{e-10}$	$5.9 \pm 0.7\text{e-10}$	$20.0 \pm 4\text{e-10}$	1.6
<i>RXTE</i>	54669.24527	1.8	$2.9^{+1.2}_{-2}$	$2.5^{+0.2}_{-0.2}$	$4.8 \pm 0.8\text{e-10}$	$5.8 \pm 0.9\text{e-10}$	$18.0 \pm 3\text{e-10}$	1.6
<i>RXTE</i>	54670.99462	3.3	$3.9^{+1.2}_{-2}$	$2.5^{+0.1}_{-0.1}$	$5.5 \pm 0.7\text{e-10}$	$6.5 \pm 0.6\text{e-10}$	$23.2 \pm 4\text{e-10}$	1.5
<i>RXTE</i>	54672.97638	1.7	$4.1^{+1.2}_{-2}$	$2.7^{+0.2}_{-0.2}$	$4.9 \pm 0.5\text{e-10}$	$5.8 \pm 0.6\text{e-10}$	$21.5 \pm 5\text{e-10}$	1.3
<i>RXTE</i>	54674.73749	1.8	$4.2^{+1.2}_{-2}$	$2.7^{+0.2}_{-0.3}$	$4.6 \pm 0.7\text{e-10}$	$5.5 \pm 0.6\text{e-10}$	$20.8 \pm 5\text{e-10}$	1.6
<i>Swift</i> ^{b,d}	54674.94054	0.1	$3.4^{+0.4}_{-0.4}$	$2.0^{+0.4}_{-0.4}$	$7.3 \pm 0.5\text{e-10}$	$8.3 \pm 0.6\text{e-10}$	$23.3 \pm 4\text{e-10}$	1.4
<i>RXTE</i>	54677.47471	3.5	$4.2^{+1.1}_{-1.2}$	$2.6^{+0.3}_{-0.2}$	$4.5 \pm 0.7\text{e-10}$	$5.3 \pm 0.8\text{e-10}$	$18.3 \pm 3\text{e-10}$	1.2
<i>RXTE</i>	54678.59119	3.3	$4.4^{+2.2}_{-2}$	$2.3^{+0.3}_{-0.1}$	$4.3 \pm 0.7\text{e-10}$	$5.0 \pm 0.8\text{e-10}$	$17.5 \pm 4\text{e-10}$	1.5
<i>RXTE</i>	54679.63915	3.2	$4.1^{+1.2}_{-2}$	$2.7^{+0.2}_{-0.3}$	$4.0 \pm 0.5\text{e-10}$	$5.1 \pm 0.6\text{e-10}$	$14.9 \pm 5\text{e-10}$	1.3
<i>RXTE</i>	54680.29712	2.1	$2.9^{+1.2}_{-1.4}$	$2.5^{+0.2}_{-0.3}$	$3.7 \pm 0.5\text{e-10}$	$4.3 \pm 0.6\text{e-10}$	$13.2 \pm 3\text{e-10}$	1.5
<i>RXTE</i>	54681.86101	3.5	$3.6^{+1.2}_{-2}$	$2.3^{+0.2}_{-0.2}$	$3.3 \pm 0.5\text{e-10}$	$3.7 \pm 0.6\text{e-10}$	$14.7 \pm 3\text{e-10}$	1.1
<i>RXTE</i>	54682.71212	7.0	$2.3^{+0.9}_{-0.9}$	$2.5^{+0.1}_{-0.1}$	$1.5 \pm 0.04\text{e-10}$	$1.9 \pm 0.05\text{e-10}$	$6.5 \pm 1.4\text{e-10}$	1.1
<i>RXTE</i>	54683.89045	3.5	$2.2^{+1.2}_{-1.2}$	$2.2^{+0.1}_{-0.2}$	$0.95 \pm 0.07\text{e-10}$	$1.2 \pm 0.5\text{e-10}$	$3.4 \pm 0.6\text{e-10}$	0.8
<i>RXTE</i>	54684.93786	1.8	$2.2^{+0.9}_{-1}$	$2.2^{+0.2}_{-0.2}$	$0.89 \pm 0.03\text{e-10}$	$1.4 \pm 0.3\text{e-10}$	$3.2 \pm 0.5\text{e-10}$	0.9
<i>RXTE</i>	54685.80434	1.6	$2.0^{+1.0}_{-0.6}$	$2.2^{+0.1}_{-0.1}$	$0.84 \pm 0.06\text{e-10}$	$1.1 \pm 0.07\text{e-10}$	$3.0 \pm 0.6\text{e-10}$	1.2
<i>RXTE</i>	54686.31211	2.5	$1.3^{+0.9}_{-0.4}$	$2.3^{+0.1}_{-0.1}$	$0.94 \pm 0.06\text{e-10}$	$1.2 \pm 0.6\text{e-10}$	$3.5 \pm 0.8\text{e-10}$	0.8
<i>RXTE</i>	54687.04156	0.7	4.0^{+2}_{-2}	$2.5^{+0.2}_{-0.2}$	$1.2 \pm 0.1\text{e-10}$	$1.5 \pm 0.2\text{e-10}$	$5.2 \pm 1.2\text{e-10}$	0.7
<i>RXTE</i>	54687.35989	2.7	$2.7^{+1.2}_{-1.2}$	$2.3^{+0.1}_{-0.1}$	$1.1 \pm 0.06\text{e-10}$	$1.5 \pm 0.06\text{e-10}$	$4.4 \pm 0.9\text{e-10}$	1.3
<i>RXTE</i>	54687.43638	2.2	$2.7^{+1.3}_{-1.3}$	$2.3^{+0.1}_{-0.1}$	$1.1 \pm 0.07\text{e-10}$	$1.4 \pm 0.08\text{e-10}$	$4.4 \pm 1.0\text{e-10}$	1.5
<i>RXTE</i>	54688.34471	2.4	$2.7^{+1.2}_{-1.2}$	$2.4^{+0.1}_{-0.1}$	$1.2 \pm 0.05\text{e-10}$	$1.6 \pm 0.05\text{e-10}$	$4.9 \pm 2\text{e-10}$	1.2
<i>RXTE</i>	54690.23786	2.2	$2.3^{+1.1}_{-1.0}$	$2.3^{+0.1}_{-0.1}$	$1.4 \pm 0.06\text{e-10}$	$1.9 \pm 0.1\text{e-10}$	$5.5 \pm 1\text{e-10}$	1.4
<i>RXTE</i>	54691.67712	1.8	$1.9^{+1.1}_{-1.0}$	$2.3^{+0.1}_{-0.1}$	$1.5 \pm 0.07\text{e-10}$	$2.0 \pm 0.07\text{e-10}$	$5.6 \pm 1.5\text{e-10}$	1.6
<i>RXTE</i>	54692.13508	1.6	1.7^{+1}_{-1}	$2.2^{+0.1}_{-0.1}$	$1.5 \pm 0.07\text{e-10}$	$2.0 \pm 0.06\text{e-10}$	$5.6 \pm 1.1\text{e-10}$	1.3
<i>RXTE</i>	54692.74785	1.2	$3.1^{+1.3}_{-1.3}$	$2.3^{+0.1}_{-0.1}$	$1.8 \pm 0.07\text{e-10}$	$2.3 \pm 0.1\text{e-10}$	$6.8 \pm 1.2\text{e-10}$	1.3
<i>RXTE</i>	54696.71397	0.5	$3.2^{+2}_{-1.5}$	$2.3^{+0.1}_{-0.1}$	$2.1 \pm 0.1\text{e-10}$	$2.8 \pm 0.2\text{e-10}$	$8.2 \pm 1.6\text{e-10}$	1.1

^a Unabsorbed flux. ^b *Swift*/XRT was operating in WT mode. ^{c,d} Intermediate long and short pre-burst persistent emission, respectively.

the different fits. In this case the best fit values and the extrapolated peak flux are acceptable values in the order of 20% which is inside our BAT extrapolation error box.

In Table 2 we report all the measured burst parameters.

Table 2. Parameters of the Intermediate long burst.

F_{peak}^a ($\text{erg cm}^{-2} \text{ s}^{-1}$)	$8.2 \pm 2 \times 10^{-8}$
f_b^c (erg cm^{-2})	$7.6 \pm 0.3 \times 10^{-6}$
$\tau \equiv f_b/F_{\text{peak}}$ (sec)	92 ± 22
$\gamma \equiv F_{\text{pers}}^c/F_{\text{peak}}$	$2.2 \pm 0.5 \times 10^{-2}$

^aUnabsorbed flux (0.1–100 keV). ^bFluence (0.1–100 keV).

^cUnabsorbed persistent flux $F_{\text{pers}} = (1.8 \pm 0.2) \times 10^{-9} \text{ erg cm}^{-2} \text{ s}^{-1}$ (0.1–100 keV).

From the time-resolved spectral results we converted the light curve count rates to flux and found also in this

case, that a double exponential function are required to fit the intermediate long burst decay. The derived e-folding time are $\tau_1 \approx 40 \pm 3 \text{ s}$ and $\tau_2 \approx 221 \pm 9 \text{ s}$, respectively with a $\chi_{red}^2 = 1.1$ (for 163 d.o.f.). In this fit we included also the first two BAT data points. Note, using only the XRT data points we found the same results within the error bars. A single exponential function is found to be inadequate with $\chi_{red}^2 = 5.2$ (166 d.o.f.). In order to compare the decay tail with the intermediate long X-ray burst from 2S 0918-549 and SLX 1737-282 (in 't Zand et al., 2005; Falanga et al., 2008), and with the decay cooling model from Cumming & Macbeth (2004) we fitted the data with a power-law, and found an index of $\Gamma = -2.14 \pm 0.02$ with a $\chi_{red}^2 = 1.7$ (167 d.o.f.). In Fig. 6 we show the double exponential best fit (upper panel) and in a log-log scale the power-law best fit (lower panel) to the data. The double exponential fit is statistically preferred over an power-law fit by the F-test probability of 1.2×10^{-14} . Similar values of $\tau_1 = 44 \pm 7 \text{ s}$ and $\tau_2 = 218 \pm 55 \text{ s}$ or power-law index $\Gamma = -1.97 \pm 0.09$ can be obtained from the fit to the




Fig. 4. *Swift*/BAT (15–35 keV) data of the intermediate long peak black-body spectra and best fit model, $kT_{\text{bb}} \approx 2.7$, (solid line). Upper limit, $kT_{\text{bb}} = 3$ keV, acceptable fit (dashed-line) and lower, $kT_{\text{bb}} = 2$, keV unacceptable fit (dotted-line).




Fig. 5. Evolution of the spectral parameters, as inferred from *Swift*/BAT (first 86 s) and *Swift*/XRT (833 s) observations. The bolometric luminosity is calculated by assuming a distance of 6.2 kpc, see Sec. 3.1. The bottom panel shows the χ^2_{red} values for these fits.

bolometric luminosity reported in Fig. 5. Also in this case the double exponential fit is statistically preferred over an power-law fit by the F-test probability of 6.6×10^{-3} .

The decay time determined with the light curve in unit of count rates, is restricted to the XRT 0.3–10 keV energy band. This decay time does not take into account the different rise and cooling effect for different energy bands during the outburst, where converting the rates to the bolometric flux or luminosity using the time-resolved spectral results

a more realistic decay time can be obtained. Therefore, in the following we consider only the $\tau_1 \approx 40$ s and $\tau_2 \approx 221$ s or the power-law index -2.14 . In Table 2 we report the $\tau \equiv f_b/F_{\text{peak}} = 92 \pm 22$ s which is only valid if one exponential function describe the decay tail. In our case this value is consistent within the error bars if we consider the double exponential decay as; $f_b/F_{\text{peak}} = \epsilon(\tau_1 - \tau_2) + \tau_2 = 71 \pm 12$, where $\epsilon = F_x/F_{\text{peak}} = 0.83$ and F_x is the peak flux for τ_1 .




Fig. 6. Top panel: We show the BAT and XRT intermediate long burst light curve with the double exponential best fit curve. Bottom panel: The same data presented in a log-log scale with the power-law best fit model.

Table 3. Fit parameters of the time-averaged BAT and XRT burst spectra.

ΔT (s)	L_{bol} ($10^{38} \text{ erg s}^{-1}$)	kT_{bb} (keV)	R_{bb} (km)	$\chi^2/\text{d.o.f.}$ –
25 ^a	2.8 ± 0.8	2.54 ± 0.33	9.3 ± 2.5	6.9/7
40 ^a	3.8 ± 0.9	2.68 ± 0.21	9.7 ± 2.5	4.9/7
22 ^a	3.4 ± 0.6	2.68 ± 0.35	10.5 ± 2.5	10.1/7
10	1.0 ± 0.1	1.83 ± 0.14	10.7 ± 1.2	54.8/51
10	0.82 ± 0.09	1.81 ± 0.14	9.9 ± 1.1	39.2/43
10	0.63 ± 0.07	1.65 ± 0.14	10.5 ± 1.3	43.4/37
10	0.58 ± 0.07	1.70 ± 0.15	9.5 ± 1.3	45.3/33
10	0.42 ± 0.05	1.46 ± 0.12	10.8 ± 1.5	38.5/28
10	0.41 ± 0.05	1.52 ± 0.14	9.9 ± 1.4	27.0/26
10	0.38 ± 0.05	1.46 ± 0.16	10.3 ± 1.8	19.4/24
10	0.33 ± 0.05	1.43 ± 0.16	9.9 ± 1.8	33.4/22
10	0.29 ± 0.04	1.36 ± 0.19	10.5 ± 2.3	26.7/19
10	0.30 ± 0.04	1.29 ± 0.13	11.8 ± 2.1	26.4/21
50	0.22 ± 0.01	1.23 ± 0.06	11.1 ± 0.9	109.4/87
50	0.17 ± 0.01	1.19 ± 0.06	10.4 ± 1.1	70.7/70
50	0.13 ± 0.01	1.08 ± 0.06	11.0 ± 1.2	50.3/59
50	0.11 ± 0.01	1.00 ± 0.06	11.7 ± 1.6	53.0/50
100	0.077 ± 0.009	0.98 ± 0.06	10.4 ± 1.4	100.1/75
100	0.054 ± 0.007	1.00 ± 0.09	8.3 ± 1.6	45.1/61
100	0.042 ± 0.007	0.96 ± 0.11	7.9 ± 1.9	60.3/52
100	0.033 ± 0.009	1.00 ± 0.16	6.4 ± 2.0	38.2/45
130	0.026 ± 0.009	0.89 ± 0.14	7.3 ± 2.4	49.4/52

^aBAT spectrum.

2.3. Short burst light curves and spectra




Fig. 7. Top panel: The short type I X-ray burst detected from XTE J1701-407 on July 27, 2008. For comparison of the rise time and duration we show also the intermediate long burst (see also Fig. 3). The *Swift*/BAT (15–150 keV) light curves are shown with a time bin of 2 s.

In Fig. 7 we show the *Swift*/BAT 15–150 keV short burst light curve. The time T_0 expressed in UTC corresponds to 22^h31^m19^s. The rise time is 1 ± 0.5 s where the total duration is ≈ 8 s. To determine the rise time and burst duration we rebinned the light curve to 1 s. For this short burst the XRT light curve starts with a delay of 180 s after the beginning of the burst as seen by BAT. In the XRT data (0.3–10 keV) the burst has not been detected, so the total duration of this burst has to be considered < 180 s. In Fig. 7 we show the short burst, and for comparison we plot also the intermediate long burst seen by BAT.

Given the short duration and statistic of the burst light curve we were not able to study a time-resolved spectral analysis. Here we report the burst spectrum analysis integrated over 6 s in order to measure as accurate as possible the peak flux. Burst spectra were well fit by a simple black-body model. The inferred BB temperature, $kT_{\text{bb}} = 3.9 \pm 0.5$ keV with an unabsorbed bolometric peak fluxes, $F_{\text{bol}} = 3.8 \pm 1.1 \times 10^{-8}$ (0.1–100 keV). The burst occurred at a flux persistent level of $2.3 \pm 0.4 \times 10^{-9}$ erg cm⁻² s⁻¹. This short burst occurred at a comparable persistent emission level as the intermediate long burst.

3. Discussion

3.1. Source distance, persistent flux, and accretion rate

When a burst undergoes a photospheric radius expansion (PRE), the source distance can be determined based on the assumption that the bolometric peak luminosity is saturated at the Eddington limit, L_{Edd} , (e.g., Lewin, van Paradijs & Taam, 1993; Kuulkers et al., 2003). During the PRE episode, while the bolometric luminosity remains constant at the Eddington value, the high energy flux may display a double peak profile and/or a delay in the rise time (e.g., Kuulkers et al., 2002; Galloway et al., 2006a; Falanga et al., 2007). The BAT light curve shows a slow rise time of ≈ 45 s, which is typically observed at high

energy in intermediate long helium bursts with PRE (e.g., Kuulkers et al., 2002; Molkov et al., 2005; Falanga et al., 2008).

Since *Swift*/XRT missed the first 133 second of the burst from XTE J1701-407 we cannot directly determine whether this burst underwent a PRE. This issue cannot be resolved with BAT time-resolved spectral analysis due to limited statistics (see Fig. 5). However, the comparison with other PRE bursts showing this slow rise time at high energy leads us to take the observed profile in the BAT light curve as an evidence for a PRE during the first ≈ 50 s of the intermediate long burst, in which case this should correspond to the timescale for the photosphere to fall back to the neutron star surface. For the short burst no conclusion can be drawn as to whether a PRE event occurred (see Fig. 7).

Another possibility is that the burst has an intrinsically long rise time. For example, the mixed H/He bursts observed from GS 1826-24 (Galloway et al., 2004) do not show PRE, but have a rise lasting for 10 seconds set by hydrogen burning (Heger et al., 2007). The duration of the bursts from GS 1826-24 is much less than the long burst from XTE J1701-407. It is not clear whether a long duration mixed H/He burst could have a 50 second rise (see discussion of light curves in section 3.5). The fact that helium burning can be extremely rapid, whereas hydrogen burning involves slow weak interactions means that the rise time is longer when hydrogen is present, and PRE much less likely. For example, Fujimoto et al. (1987) derived a critical helium fraction necessary to achieve PRE.

Assuming a bolometric peak luminosity equal to the Eddington value for a He type I X-ray burst ($L_{\text{Edd}} \approx 3.8 \times 10^{38}$ ergs⁻¹, as empirically derived by Kuulkers et al., 2003), we obtain the source distance of $d = 6.2^{+1.6}_{-0.9}$ kpc. For comparison, the theoretical value of this distance (e.g., Lewin, van Paradijs & Taam, 1993) found by assuming a He atmosphere and canonical NS parameters (1.4 solar mass and radius of 10 km), is $5.5^{+1.3}_{-0.8}$ kpc. Note that the source could be closer if the peak luminosity of the burst was lower than the pure helium Eddington limit. For example, we cannot rule out that the burst did not show PRE, in which case the peak luminosity could have been sub-Eddington. Alternatively, taking the peak luminosity to be the Eddington luminosity for solar composition ($X_0 = 0.7$) gives a distance of ≈ 4 kpc. In the following we consider $d \approx 6.2$ kpc as a fiducial distance, and comment on how our conclusions would change if the source was closer.

The best fit to the 2–20 keV persistent emission spectrum of XTE J1701-407 required an absorbed simple power-law model with $\Gamma \approx 2.1$. Assuming a distance of 6.2 kpc, the estimated intermediate long burst pre-burst persistent unabsorbed flux between 0.1–100 keV, $F_{\text{pers}} \approx 1.9 \times 10^{-9}$ erg cm⁻² s⁻¹, translates into a bolometric luminosity of $L_{\text{pers}} \approx 8.3 \times 10^{36}$ erg s⁻¹, or $\approx 2.2\% L_{\text{Edd}}$.

The local accretion rate per unit area is then given by $L_{\text{pers}} = 4\pi R^2 \dot{m} (GM/R)/(1+z)$, or

$$\dot{m} = 4.0 \times 10^3 \text{ g cm}^{-2} \text{ s}^{-1} \left(\frac{R}{11.2 \text{ km}} \right)^{-1} \left(\frac{M}{1.4 M_{\odot}} \right)^{-1} \left(\frac{d}{6.2 \text{ kpc}} \right)^2 \left(\frac{1+z}{1.26} \right) (1)$$

A convenient unit for accretion rate is the Eddington accretion rate. Here, we define the local Eddington accretion rate as $\dot{m}_{\text{Edd}} \equiv 1.8 \times 10^5 \text{ g cm}^{-2} \text{ s}^{-1}$, the local accretion rate onto a $M = 1.4 M_{\odot}$, and $R = 11.2$ km neutron

star that gives an accretion luminosity equal to the empirically derived Eddington luminosity $3.8 \times 10^{38} \text{ erg s}^{-1}$ from Kuulkers et al. (2003). This gives $\dot{m}/\dot{m}_{\text{Edd}} = 2.2\%$.

3.2. The energy, ignition depth, and recurrence time of the long burst

The observed energy of the long burst allows us to estimate the ignition depth. The measured fluence of the burst is $f_b = 7.6 \times 10^{-6} \text{ erg cm}^{-2}$, giving a burst energy release $E_{\text{burst}} = 4\pi d^2 f_b = 3.5 \times 10^{40} \text{ ergs}$ ($d/6.2 \text{ kpc}$)². The ignition depth is given by $E_{\text{burst}} = 4\pi R^2 y_{\text{ign}} Q_{\text{nuc}}/(1+z)$, or

$$y_{\text{ign}} = 1.8 \times 10^9 \text{ g cm}^{-2} \left(\frac{d}{6.2 \text{ kpc}} \right)^2 \left(\frac{Q_{\text{nuc}}}{1.6 \text{ MeV/nucleon}} \right)^{-1} \left(\frac{R}{11.2 \text{ km}} \right)^{-2} \left(\frac{1+z}{1.26} \right) \quad (2)$$

The value of $Q_{\text{nuc}} \approx 1.6 \text{ MeV}$ corresponds to the nuclear energy release per nucleon for complete burning of helium to iron group elements. Including hydrogen with mass-weighted mean mass fraction $\langle X \rangle$ gives $Q_{\text{nuc}} \approx 1.6 + 4\langle X \rangle \text{ MeV/nucleon}$ (Galloway et al., 2004), where we include losses due to neutrino emission following Fujimoto et al. (1987). For $\langle X \rangle = 0.7$, the solar composition value, $Q_{\text{nuc}} = 4.4 \text{ MeV/nucleon}$, and $y_{\text{ign}} = 6.5 \times 10^8 \text{ g cm}^{-2}$.

At an accretion rate of $4 \times 10^3 \text{ g cm}^{-2} \text{ s}^{-1}$, the recurrence time corresponding to a column depth of $y_{\text{ign}} = 1.8 \times 10^9 \text{ g cm}^{-2}$ (pure helium composition) is $\Delta t = (y_{\text{ign}}/\dot{m})(1+z) = 6.6 \text{ days}$, or for $y_{\text{ign}} = 6.5 \times 10^8 \text{ g cm}^{-2}$ (solar composition) is $\Delta t = 2.4 \text{ days}$. These derived recurrence times are independent of the assumed distance. The intermediate long burst, as the first observed burst, occurred 40 days after the detection of XTE J1701-407. The effective exposure time on the source from the beginning of the outburst to the intermediate long burst was 0.34 days, and 0.14 days between both bursts. The elapsed time on the source is thus too short compared to the theoretically derived recurrence time to allow us to get an observational measurement of the recurrence time.

3.3. Theoretical comparison with ignition models

To try to understand the nuclear burning processes responsible for the long burst, we compare the observed properties with type I X-ray burst ignition models. The ignition conditions are calculated as described in Cumming & Bildsten (2000), but we take $R = 11.2 \text{ km}$, $M = 1.4 M_{\odot}$, and the energy release in hot CNO burning⁴ $E_H = 6.0 \times 10^{18} \text{ erg g}^{-1}$. The results are shown in Table 4 for several different choices of accreted hydrogen fraction X_0 , accretion rate $\dot{m}$ and flux from the crust Q_b . The ignition conditions are calculated by modeling the temperature profile of the accumulating fuel layer, and adjusting the thickness of the layer until a condition for thermal runaway is met at the base. These models do not include the effects of previous bursts on the ignition conditions (e.g. Woosley et al. 2004), and we have not included gravitational sedimentation which is important at low accretion rates (Peng et al., 2007). In the following, we

consider three possibilities: 1) accretion of pure helium, 2) hydrogen-rich matter at solar, and 3) low metallicity.

3.3.1. Pure helium accretion

For pure helium accretion, the accumulating fuel layer is heated from below by a flux $\dot{m}Q_b$ from the crust, where Q_b is the energy per accreted nucleon flowing outwards from the crust. We set the accretion rate at the observed value $\dot{m}/\dot{m}_{\text{Edd}} = 2.2\%$, and adjust the value of Q_b until we find ignition at the inferred column for pure helium burning $y_{\text{ign}} = 1.8 \times 10^9 \text{ g cm}^{-2}$ (and therefore obtain the observed burst energy). A flux from below $Q_b = 0.5 \text{ MeV/nucleon}$ matches the observed burst energy at 2.2% Eddington. We also include models with $Q_b = 0.3$ and 0.7 MeV/nucleon in Table 4 to show the sensitivity of the ignition depth to the amount of heat from below. Note that since the combination $\dot{m}Q_b$ sets the total flux heating the layer, an increase or decrease in Q_b keeping $\dot{m}$ fixed is equivalent to increasing or decreasing $\dot{m}$ with Q_b fixed. Therefore, if the inferred accretion rate is smaller by some factor, the value of Q_b needed to match the observed energy will be larger by the same factor. Similarly, a source distance less than 6.2 kpc can be accommodated by increasing Q_b .

The outwards flux expected from the crust depends on accretion rate and the core neutrino emissivity (Brown, 2000), ranging from $\dot{m}Q_b \approx 0.1 \text{ MeV}$ per nucleon at high accretion rates up to $\dot{m}Q_b \approx 1.5 \text{ MeV}$ per nucleon at low accretion rates. The value depends on how much of the total $\approx 1.5 \text{ MeV}$ per nucleon heat released in the crust by pycnonuclear and electron capture reactions (Haensel & Zdunik, 1990, 2008) is conducted into the core and released as neutrinos compared to being conducted outwards. The number of 0.5 MeV/nucleon that we infer here is quite reasonable for an accretion rate of 2.2% Eddington; for example, the models calculated by Cumming et al. (2006) for persistently accreting sources have $Q_b = 0.3$ to 0.9 MeV/nucleon at this accretion rate. Galloway & Cumming (2006b) modelled the X-ray bursts observed from SAX J1808.4-3658 and found $Q_b \approx 0.3 \text{ MeV/nucleon}$ for $\dot{m} \approx 3\% \dot{m}_{\text{Edd}}$.

3.3.2. Accretion of hydrogen rich matter with solar metallicity

Next, we consider that the source is accreting hydrogen rich matter with the solar hydrogen fraction $X_0 = 0.7$ and a metallicity similar to solar, with mass fraction of CNO elements $Z_{\text{CNO}} = 0.02$. In that case, the hydrogen burns at a fixed rate during accumulation of the fuel layer, by the beta-limited hot CNO cycle of Hoyle & Fowler (1965). The hydrogen depletes at a column depth

$$y_d = 1.5 \times 10^8 \text{ g cm}^{-2} \left(\frac{\dot{m}}{0.022\dot{m}_{\text{Edd}}} \right) \left(\frac{Z_{\text{CNO}}}{0.02} \right)^{-1} \left(\frac{X_0}{0.7} \right) \quad (3)$$

smaller than the inferred ignition depth, so that a thick layer of pure helium accumulates and ignites beneath a steady hydrogen burning shell. Therefore, even though solar composition material is accreted, the mean hydrogen fraction at ignition in the fuel layer is very small, giving Q_{nuc} close to the value 1.6 MeV/nucleon for pure helium.

For a CNO mass fraction $Z_{\text{CNO}} = 0.02$, the observed burst energy is obtained for an accretion rate three times

⁴ Here we adopt the value of E_H from Wallace & Woosley (1981), which includes neutrino losses. These were not included by Cumming & Bildsten (2000).

Table 4. Type I X-ray burst ignition conditions^a

Model	$\dot{m}^b$ (% $\dot{m}_{\text{Edd}}$)	Z_{CNO}	X_0^c	Q_b	$y_{\text{ign},9}$	$T_{\text{ign},8}$	$\langle X \rangle$	X_b	Q_{nuc}	E_{40}	Δt (days)
Pure helium accretion ^d											
1	2.2	0.02	0	0.5	1.8	1.4	0	0	1.6	3.4	6.5
2	2.2	0.02	0	0.3	7.4	1.2	0	0	1.6	14	27
3	2.2	0.02	0	0.7	0.84	1.5	0	0	1.6	1.6	3.1
Accretion hydrogen rich material											
4	2.2	0.02	0.7	0.5	0.13	2.0	0.38	0.07	3.1	0.49	0.48
5	0.69	0.02	0.7	0.5	1.6	1.4	0.01	0	1.64	3.2	19
6	2.2	0.001	0.7	0.1	0.67	1.7	0.62	0.54	4.1	3.3	2.5
7	2.2	0.001	0.7	0.5	0.53	1.8	0.63	0.43	4.1	2.7	2.0

^a Models 1, 5, and 6 provide a good match to the observed burst energy of 3.5×10^{40} ergs. In addition, models 1 and 6 have an accretion rate that matches the value inferred from the persistent luminosity.

^b We define $\dot{m}_{\text{Edd}} = 1.8 \times 10^5 \text{ g cm}^{-2} \text{ s}^{-1}$, the local accretion rate onto a $1.4 M_{\odot}$, $R = 11.2 \text{ km}$ neutron star that gives an accretion luminosity equal to the empirically derived Eddington luminosity $3.8 \times 10^{38} \text{ erg s}^{-1}$ from Kuulkers et al. (2003).

^c The hydrogen mass fractions are: in the accreted material X_0 , at the base of the layer at ignition X_b , and the mass-weighted mean value in the layer at ignition $\langle X \rangle$.

^d Note that the ignition conditions for pure helium accretion do not depend on the choice of Z_{CNO} .

lower than inferred from the observed X-ray luminosity $\dot{m}/\dot{m}_{\text{Edd}} = 0.69\%$ (model 5 in Table 4). At an accretion rate of 2.2% Eddington, the burst energy is a factor of seven too small (model 4). Reducing the distance to the source does not help because it would change both the inferred $\dot{m}$ and burst energy by the same factor.

3.3.3. Accretion of hydrogen rich matter with low metallicity

In the third scenario, we consider that the material is hydrogen rich, but with a low metallicity. Model 6 has a burst energy close to the observed energy at the inferred accretion rate of 2.2% Eddington, with a CNO mass fraction $Z = 10^{-3}$, roughly 10% of the solar metallicity. In this case, the amount of hot CNO burning is reduced substantially, so that hydrogen permeates the entire fuel layer at ignition. Hot CNO burning still operates at a low level and causes some preheating of the fuel layer. The hydrogen increases the amount of nuclear release during the burst, giving $Q_{\text{nuc}} \approx 4 \text{ MeV/nucleon}$, and an ignition depth three times smaller than for the pure helium case. The low metallicity ignition is less sensitive to Q_b than the pure helium ignition. Model 7, which has the same conditions as model 6, but with $Q_b = 0.5$ rather than $Q_b = 0.1$, has a burst energy within almost 20% of the observed value. Similar to the pure helium ignitions, a closer distance can be accommodated by varying Q_b .

One caveat regarding the low metallicity ignition models is that there can be substantial heating of the accumulating fuel layer because of nuclear reactions associated with the ashes of previous bursts. Woosley et al. (2004) found that the burst behavior at accretion rates ≈ 0.1 Eddington was insensitive to the metallicity of the accreted material due to this effect. However, they found that at lower accretion rates there was good agreement with the ignition models presented here (see Woosley et al., 2004, Table 9).

3.3.4. Summary of ignition models

We present three ignition models in Table 4 that fit the observations. Models 1 and 6 have the correct burst energy and accretion rate. They correspond to accretion of pure helium (model 1), for which the layer is heated by the outwards flux from the neutron star crust, and for accretion of hydrogen rich material with low metallicity (model 6), for which a low level of hydrogen burning preheats the layer during accumulation, but the hydrogen fraction at ignition is substantial and makes a significant contribution to the burst energetics. Third, model 5 has the correct burst energy, but at an accretion rate three times lower than observed. Given the uncertainties in translating the observed X-ray luminosity to accretion rate, it seems worthwhile considering this model further. In this model, the accreted composition is hydrogen-rich with a solar metallicity. This leads to depletion of the hydrogen by the hot CNO cycle and the build-up of a thick layer of pure helium beneath the hydrogen shell.

There is a fourth possibility, which is that the source is accreting hydrogen-rich material, but the hydrogen burns unstably in a series of short flashes. The helium produced in the short flashes builds up and make a pure helium layer that ignites to give the long burst. This is similar to model 5, but with unstable rather than stable hydrogen burning.

One way to distinguish the various possible explanations for the burst energetics would be a recurrence time measurement: as shown in Table 4, the different scenarios predict differences in recurrence time. In fact, this is equivalent to a measurement of the α parameter⁵ for the bursts, which would indicate the fuel type (e.g. $\alpha \approx 40$ for solar hydrogen abundance as found for example in GS 1826-24, Galloway et al. 2004).

While this paper was in preparation, a similar analysis of the intermediate burst from XTE J1701-407 was carried out and reported in the preprint by Linares et al. (2008b). The bolometric peak flux for the long burst and therefore

⁵ The quantity α is defined as the ratio of the total energy emitted in the persistent flux to that emitted in a burst, $\alpha = F_{\text{pers}} \Delta t / f_b$, where Δt is the time interval between two bursts.

the limits on the source distance are consistent with the values we obtain here. However, the persistent luminosity and burst energy given in Linares et al. (2008b) are a factor of two smaller than our values, being restricted to the energy range 1–50 keV, whereas we have included a bolometric correction. These values are used in the interpretation of the long burst, resulting in an accretion rate of $\dot{m} = 1.9 \times 10^3 \text{ g cm}^{-2} \text{ s}^{-1} = 1.1\% \dot{m}_{\text{Edd}}$ and a burst energy of $E_b = 1.6 \times 10^{40} \text{ ergs}$, both a factor of two smaller than the values we find in this paper. Repeating the ignition calculations presented earlier for this lower accretion rate, we find that accretion of solar composition material ($X_0 = 0.7$) gives a burst energy of $E_b = 1.5 \times 10^{40} \text{ ergs}$ for $Q_b = 0.1 \text{ MeV/nucleon}$, and $E_b = 1.0 \times 10^{40} \text{ ergs}$ for $Q_b = 0.5 \text{ MeV/nucleon}$. Therefore, accretion of solar composition naturally explains the burst energetics for these values of E_b and $\dot{m}$. For pure helium accretion at this rate, we find that $Q_b = 1.5 \text{ MeV/nucleon}$ is required to achieve a burst energy of $1.5 \times 10^{40} \text{ ergs}$ (lower values of Q_b result in a deeper ignition and more energetic burst). Whereas this is a larger value than expected at this accretion rate, it is within the range of the total energy released in the crust (Haensel & Zdunik, 2008). In addition, since the flux heating the layer is $\propto \dot{m}Q_b$, the requirements on Q_b can be relaxed if the true accretion rate is larger than assumed here. Therefore, we find that the energetics argument given in Linares et al. (2008b) against explaining this burst as pure helium is overstated.

3.4. Constraints from the light curve

The shape and duration of the light curve offer another way to determine the composition of the fuel that burns during the burst. While we do not have models available with exactly the same burst energy, the low accretion rate models from Woosley et al. (2004) are within a factor of two to four in energy and accretion rate, and so we compare these models with the observed light curve. These models have $\dot{M} = 3.5 \times 10^{-10} M_{\odot} \text{ yr}^{-1}$, corresponding to a local accretion rate $\dot{m} = 1\% \dot{m}_{\text{Edd}}$.

First, we compare the observed light curve with burst 2 from model zm of Woosley et al. (2004) (solid curve in Fig. 8). This model is for accretion of hydrogen rich ($X_0 \approx 0.7$) matter with a low metallicity, $Z = 10^{-3}$, and so is similar to model 6 in Table 4. The burst has a total energy release of $2.0 \times 10^{40} \text{ ergs}$, just less than a factor of two smaller than the observed burst. The recurrence time is 3.0 days, and ignition column $y_{\text{ign}} = 3.6 \times 10^8 \text{ g cm}^{-2}$. The ignition column is just less than a factor of two smaller than the observed burst. The peak luminosity of this burst is close to the Eddington luminosity for solar composition, suggesting that the distance may be closer than the 6.2 kpc assumed here. A closer distance would bring these light curves into better agreement. The model light curve has a steep decline at late times, somewhat steeper than the observed decline. An extra factor of two in ignition column would give the correct burst energy and lengthen the model burst light curve, bringing it into better agreement with the observed burst. Another point to note is that this burst has a slow rise time, lasting for several seconds, as expected for a low helium mass fraction (Fujimoto et al., 1987).

The dashed curve in Fig. 8 shows burst 3 from model Zm of Woosley et al. (2004). This model has hydrogen-rich

matter with solar metallicity, $Z = 0.02$, and so is similar to model 5 of Table 4. The hydrogen burns away in a thin shell, leaving a pure helium layer that ignites. This burst has a total energy release of $8.0 \times 10^{39} \text{ ergs}$, a factor of 4 smaller than observed. The recurrence time is 4.5 days, and ignition column depth $5.6 \times 10^8 \text{ g cm}^{-2}$. The burst reaches close to the pure helium Eddington luminosity. The duration of the burst is shorter than the observed burst by a factor of ≈ 5 –10. The slope of the decay is shallower than the observed slope. The rise time of this burst is very fast, a fraction of a second, in contrast to the much slower rise of the hydrogen-rich burst (model zm; solid line in Fig. 8).

We also include some cooling models calculated following Cumming & Macbeth (2004). For a given ignition column, an energy release per gram of 1.6 MeV per nucleon is deposited in the layer, as would be appropriate if the helium burned to iron group elements at each depth at the start of the burst. We have also computed models for a lighter ash and correspondingly smaller energy deposition (Woosley et al. 2004 find that the burning does not go all the way to iron group in their model zM), but the differences are small, and this does not change our conclusions. The cooling of the layer is then followed, with the flux from the surface limited to the Eddington luminosity for pure helium⁶. We show two examples in Fig. 8. The first has $y_{\text{ign}} = 6 \times 10^8 \text{ g cm}^{-2}$ to match the Woosley et al. (2004) model Zm burst. The second has $y_{\text{ign}} = 2 \times 10^9 \text{ g cm}^{-2}$ as needed to get the observed burst energy (models 1 and 5 in Table 4). The shape of the cooling models agrees well with the model Zm light curve, and agrees within a factor of two in the cooling timescale. Even allowing for this factor of two in the $y_{\text{ign}} = 2 \times 10^9 \text{ g cm}^{-2}$ model, the cooling is faster than the observed light curve.

In summary, although the models from Woosley et al. (2004) are not at exactly the same ignition conditions as implied by the observations of XTE J1701-407, our comparison suggests that pure helium ignition at the inferred ignition column depth (with or without a small overlying hydrogen burning shell) have cooling times that are shorter than the observed light curve. On the other hand, a hydrogen-rich composition throughout the layer, as expected for low metallicity, produces a longer lasting light curve more consistent with the observed cooling time. This conclusion does not depend on the assumed distance to the source. Both helium and hydrogen-rich burst models reach the Eddington luminosity (either the pure helium or solar composition Eddington luminosity respectively), and therefore could explain the PRE suggested by the similarity between the observed BAT light curve and other intermediate long bursts (see section 3.1). If the slow rise time is intrinsic to the burst and not due to PRE, the hydrogen rich model is preferred as the presence of hydrogen leads to a much slower rise time than for pure helium (Fig. 8). The double exponential nature of the decay may also argue for hydrogen burning during the burst. The burst profiles from GS 1826-24 (Galloway et al., 2004) are well-fit by a double exponential decay. Further modeling is required to

⁶ As noted by Woosley et al. (2004), the shape of the light curve as the luminosity begins to drop below the Eddington luminosity is likely not accurately reproduced by these models, which do not follow the outer layers in detail. We will improve our treatment of this in future work. However, we expect that the late time cooling is not sensitive to this.

study the expected burst profiles of hydrogen-rich bursts produced as a result of low metallicity accretion.




Fig. 8. Model light curves compared with the observed light curve. The solid curve shows burst 2 from model zm of Woosley et al. (2004). This burst has an ignition column and energy a factor of 2 smaller than the observed burst. The dashed curve shows burst 3 from model Zm of Woosley et al. (2004). This burst has an ignition column a factor of three to four times smaller than the observed burst. The observed bolometric flux has been converted to luminosity using a distance of 6.2 kpc. Redshift corrections have been applied to the theoretical light curves, with $1 + z = 1.26$. The dotted curves show two cooling models calculated following Cumming & Macbeth (2004), for (left to right) $y_{\text{ign}} = 6 \times 10^8 \text{ g cm}^{-2}$ corresponding to Woosley et al. (2004) model Zm (dashed curve), and $y_{\text{ign}} = 2 \times 10^9 \text{ g cm}^{-2}$ as needed to explain the observed burst energy (models 1 and 5 in Table 4).

3.5. Origin of the short burst

We have focused on the long duration burst, which has a well-measured fluence and therefore energy. For the short burst, we can only place an upper limit on its fluence. At low accretion rates, unstable ignition of hydrogen can give rise to short duration bursts (e.g., Strohmayer & Bildsten, 2006; Chenevez et al., 2007). Linares et al. (2008b) suggest that this is the origin of the short burst from XTE J1701-407, and that either (i) hydrogen-ignited short bursts are producing the helium fuel for the long burst, or that (ii) the source is accreting close to the boundary between unstable and stable hydrogen burning, and stable hydrogen burning produces the helium for the long burst. The comparison to ignition models and model light curves that we have made earlier suggest that hydrogen survives to the ignition depth, implying a low metallicity in the accreted layer. Unfortunately, this would presumably make a thermal instability driven by CNO burning less likely.

For the low metallicity model or for pure helium accretion, another explanation is that the local accretion rate was higher at the time of the short burst. The ignition depth is very sensitive to the base flux or equivalently to the product $\dot{m}Q_b$ (see for example Fig. 8 of in 't Zand et al. 2005). The persistent flux at the time of the short burst from

XTE J1701-407 was slightly larger than at the time of the long burst, even at 30%, not enough to cause a significant reduction in the ignition column depth. We note that short bursts were observed from the intermediate long X-ray burster 2S 0918-549, a suspected ultracompact binary and therefore accreting hydrogen-deficient matter. Therefore, it is not clear to us that the observation of the short burst rules out pure helium accretion in XTE J1701-407, as argued by Linares et al. (2008b).

4. Conclusions

We have compared the observed properties of the long duration burst from XTE J1701-407 with models of type I X-ray burst ignition conditions and light curves. We showed that the observed burst energy could be understood as (i) pure helium ignition, either as a result of pure helium accretion or of depletion of hydrogen by steady burning during accumulation, or (ii) as ignition of a thick layer of hydrogen-rich material with low metallicity. Comparing with model light curves, we find that the pure helium ignitions cool faster than observed. On the other hand, a hydrogen rich layer gives a longer duration light curve with a steep decline in the tail of the burst, better matching the observed light curve. Therefore we suggest that the intermediate long burst from XTE J1701-407 was powered by unstable burning of a thick layer of hydrogen rich matter with low metallicity. Long X-ray bursts caused by pure helium ignitions beneath a hydrogen shell have been identified, as for example by Galloway & Cumming (2006b) who argued that this was happening in SAX J1808.4-3658. But to our knowledge XTE J1701-407 would be the first example of a source that shows long bursts driven by a thick layer of hydrogen-rich material. The bursts from GS 1826-24 are believed to be powered by rp-process hydrogen burning, giving long ≈ 100 second tails, but the ignition depth in those bursts is an order of magnitude smaller than inferred for the long burst from XTE J1701-407, so that hydrogen can survive until helium ignition, even for solar metallicity. At the low accretion rate in XTE J1701-407, this is not the case: low metallicity is required to reduce the rate of hot CNO burning and allow hydrogen to survive until helium ignites. This implies either that this source is a burster accreting low metallicity H-rich material at a low rate, or another possibility is that heavy elements are able to sediment out from the accumulating layer at this low accretion rate (Peng et al., 2007), reducing its effective metallicity. If so, future studies of this source could be used to test the physics of sedimentation at low accretion rates.

Acknowledgements. We thank Alexander Heger for providing the burst light curves from Woosley et al. (2004) shown in Fig. 7. MF acknowledges the French Space Agency (CNES) for financial support. JC acknowledges financial support from ESA-PRODEX, Nr. 90057, and EB acknowledges ASI and MIUR. AC acknowledges support from the National Sciences and Engineering Research Council of Canada (NSERC), Le Fonds Québécois de la Recherche sur la Nature et les Technologies (FQRNT), and the Canadian Institute for Advanced Research (CIFAR). AC is an Alfred P. Sloan Research Fellow.

References

- Barthelmy, S. D. et al. 2008, GCN Circular, 7985
- Brown, E. F. 2000, ApJ, 531, 988
- Chelovekov, I. V., Grebenev, S. A. & Sunyaev, R. A. 2006, AstL, 32, 456

- Chenevez, J., Falanga, M., Brandt, S. et al. 2006, *A&A*, 449, L5
- Chenevez, J., Falanga, M., Kuulkers, E. et al. 2007, *A&A*, 469, L27
- Cornelisse, R., in 't Zand, J. J. M., Verbunt, F. et al. 2003, *A&A*, 405, 1033
- Cumming, A., & Bildsten, L. 2000, *ApJ*, 544, 453
- Cumming, A., & Bildsten, L. 2001, *ApJ*, 559, L127
- Cumming, A., & Macbeth, J. 2004, *ApJ*, 603, L37
- Cumming, A., Macbeth, J., in 't Zand, J.J.M., Page, D. 2006, *ApJ*, 646, 429
- Degenaar, N. & Wijnands, R. 2008, *Astr. Tel.*, 1572
- Falanga, M., Poutanen, J., Bonning, E. W. et al. 2007, *A&A*, 464, 1069
- Falanga, M., Chenevez, J., Cumming, A. et al. 2008, *A&A*, 484, 43
- Fujimoto, M. Y., Sztajno, M., Lewin, W. H. G., van Paradijs, J., 1987, *ApJ*, 319, 902
- Galloway, D. K., Cumming, A., Kuulkers, E. et al. 2004, *ApJ*, 601, 466
- Galloway, D. K., Muno, M.P., Hartman, J.M, et al. 2006a, *ApJS*, submitted, [arXiv:astro-ph/0608259]
- Galloway, D. K., & Cumming, A. 2006b, *ApJ*, 652, 559
- Hansen, C. J. & Van Horn, H. M., 1975, *ApJ*, 195, 735
- Haensel, P., & Zdunik, J. L. 1990, *A&A*, 227, 431
- Haensel, P., & Zdunik, J. L. 2008, *A&A*, 480, 459
- Hasinger, G. & van der Klis, M. 1989, *A&A*, 225, 79
- Heger, A., Cumming, A., Galloway, D. K., & Woosley, S. E. 2007, *ApJ*, 671, L141
- Hoyle, R., & Fowler, W. A. 1965, in *Quasi-Stellar Sources and Gravitational Collapse*, ed. I. Robinson, A. Schild, & E. L. Shucking (Chicago : Univ. Chicago Press), 17
- in't Zand, J. J. M., Cornelisse, R. & Cumming, A. *A&A*, 2004, 426, 257
- in 't Zand, J. J. M., et al. 2005, *A&A*, 441, 675
- Jahoda, K., Swank, J. H., Giles, A. B., et al., 1996, *Proc. SPIE*, 2808, 59
- Kaplan, D. & Chakrabarty, D. 2008, *Astr. Tel.*, 1630
- Kuulkers, E., et al. 2003, *A&A*, 399, 663
- Kuulkers, E. 2004, *Nucl. Phys. B*, 132, 466
- Kuulkers, E., Homan, J., van der Klis, M., et al. 2002, *A&A*, 382, 947
- Lewin, W. H. G. L., van Paradijs, J. & Taam, R. 1993, *Space Science Reviews*, 62, 223
- Linares, M. et al. 2008a, *Astr. Tel.*, 1618
- Linares, M. et al. 2008b, *MNRAS*, in press, [arXiv:astro-ph/0808.3950]
- Markwardt, C. B., Pereira, D., & Swank, J. H. 2008a, *Astr. Tel.*, 1569
- Markwardt, C. B., Cummings, J., & Krimm, H. 2008b, *Astr. Tel.*, 1616
- Molkov, S., Revnivtsev, M., Lutovinov, A. & Sunyaev, R. A. 2005, *A&A*, 434, 1069
- Peng, F., Brown, E. F., & Truran, J. W. 2007, *ApJ*, 654, 1022
- Rothschild, R. E., Blanco, P. R., Gruber, D. E., et al. 1998, *ApJ*, 496, 538
- Sakamoto, T. et al. 2008, *GCN circular*, 8034
- Starling, R. & Evans, P. 2008, *Astr. Tel.*, 1621
- Strohmayer, T. E., & Bildsten, L. 2006, in *Compact stellar X-ray sources*, ed. W. H. G. Lewin & M. van der Klis, (Cambridge: Cambridge University Press)
- Strohmayer, T. E., Markwardt, C. B. & Swank, J. H. 2008, *Astr. Tel.*, 1635
- Swank, J. & Markwardt, C. 2001, in *ASP Conf. Ser. 251, New Century of X-ray Astronomy*, ed. H. Inoue & H. Kunieda (San Francisco: ASP), 94
- Wallace, R. K. & Woosley, S. E. 1981, *ApJS*, 45, 1389
- Woosley, S. E., Heger, A., Cumming, A. et al. in 2004, *ApJS*, 151, 75